COPY OF ALL CLAIMS

1. (currently amended) A cyclohexenonequinolinoyl derivative of the formula I

$$R^4$$
 R^2
 R^1

where:

 $R^1 \qquad \text{is hydrogen, nitro, halogen, cyano, C_1-C_6-alkyl, C_1-C_6-haloalkyl, C_1-C_6-alkoxy, C_1-C_6-haloalkoxy, C_1-C_6-alkylthio, C_1-C_6-haloalkylthio, C_1-C_6-alkylsulfinyl, C_1-C_6-haloalkylsulfinyl, C_1-C_6-alkylsulfonyl, aminosulfonyl, N-$($C_1$-$C_6$-alkylsulfonyl, aminosulfonyl, N-$($C_1$-$C_6$-alkylsulfonyl), C_1-C_6-alkylsulfonyl, C_1-C

N, N-di-(C₁-C₆-alkyl) aminosulfonyl,

N-(C₁-C₆-alkylsulfonyl)amino,

N-(C₁-C₆-haloalkylsulfonyl)amino,

 $N-(C_1-C_6-alkyl)-N-(C_1-C_6-alkylsulfonyl)$ amino,

 $N-(C_1-C_6--alkyl)-N-(C_1-C_6-haloalkylsulfonyl)amino,$

phenoxy, heterocyclyloxy, or phenylthio or heterocyclylthio, it being possible for the four last-mentioned radicals to be partially or fully halogenated and/or to carry one to two of the following one to three of the following substituents:

nitro, cyano, C_1 - C_4 -alkyl, C_1 - C_4 -haloalkyl, C_1 - C_4 -alkoxy or C_1 - C_4 -haloalkoxy;

 R^2 , R^3 are hydrogen, C_1 - C_6 -alkyl, C_1 - C_6 -haloalkyl or halogen;

R⁴ is a compound IIa or IIb

$$(\mathsf{R}^6)_{\mathsf{I}} \qquad (\mathsf{R}^6)_{\mathsf{I}} \qquad \mathsf{IIb}$$

where

is halogen, OR⁷, SR⁷, SOR⁸, SO₂R⁸, OSO₂R⁸, POR⁸R⁹,

OPR⁸R⁹, OPOR⁸R⁹, OPSR⁸R⁹, NR¹⁰R¹¹, ONR¹¹R¹², N-linked

heterocyclyl or O-(N-linked heterocyclyl), it being possible for the heterocyclyl radical of the two last-mentioned substituents to be partially or fully halogenated and/or to

Ŷ

carry one to three of the following radicals:

nitro, cyano, C_1 - C_4 -alkyl, C_1 - C_4 -haloalkyl, C_1 - C_4 -alkoxy or C_1 - C_4 -haloalkoxy;

R⁶ is nitro, halogen, cyano, C_1 - C_6 -alkyl, C_1 - C_6 -haloalkyl, di- $(C_1$ - C_6 -alkoxy)methyl, $di-(C_1$ - C_6 -alkylthio)methyl, $(C_1$ - C_6 -alkoxy)(C_1 - C_6 -alkylthio)methyl, hydroxyl, C_1 - C_6 -alkoxy, C_1 - C_6 -haloalkoxy, C_1 - C_6 -alkoxycarbonyloxy, C_1 - C_6 -alkylthio, C_1 - C_6 -alkylthio, C_1 - C_6 -alkylthio, C_1 - C_6 -alkylsulfinyl,

C₁-C₆-haloalkylsulfinyl, C₁-C₆-alkylsulfonyl,

C₁-C₆-haloalkylsulfonyl, C₁-C₆-alkylcarbonyl,

C₁-C₆-haloalkylcarbonyl, C₁-C₆-alkoxycarbonyl or

C₁-C₆-haloalkoxycarbonyl;

or

two radicals , which are linked to the same carbon, together form an -O-(${\rm CH_2}$)_m-O-, -O-(${\rm CH_2}$)_m-S-, -S-(${\rm CH_2}$)_m-S-, -O-(${\rm CH_2}$)_n- or -S-(${\rm CH_2}$)_n chain which is unsubstituted or substituted by one to three radicals from the following group: halogen, cyano, ${\rm C_1}$ - ${\rm C_4}$ -alkyl, ${\rm C_1}$ - ${\rm C_4}$ -haloalkyl or ${\rm C_1}$ - ${\rm C_4}$ -alkoxycarbonyl;

or

two radicals , which are linked to the same carbon, together form a -(CH_2)_p chain which possibly is interrupted by oxygen or sulfur and/or is unsubstituted or substituted by one to four radicals from the following group: halogen, cyano, C_1 - C_4 -alkyl, C_1 - C_4 -haloalkyl or C_1 - C_4 -alkoxycarbonyl;

or

two radicals , which are linked to the same carbon, together form a methylidene group which is unsubstituted or substituted by one or two radicals from the following group: halogen, hydroxyl, formyl, cyano, C_1 - C_6 -alkyl, C_1 - C_6 -haloalkyl, C_1 - C_6 -alkoxy, C_1 - C_6 -haloalkoxy, C_1 - C_6 -alkylthio, C_1 - C_6 -haloalkylthio, C_1 - C_6 -alkylsulfinyl, C_1 - C_6 -haloalkylsulfonyl or C_1 - C_6 -haloalkylsulfonyl;

or

two radicals, which are linked to the same carbon, together with this carbon form a carbonyl group;

or

two radicals, which are linked to different carbons, together form a $-(CH_2)_n$ chain which is unsubstituted or

substituted by one to three radicals from the following group: halogen, C_1 - C_6 -alkyl, C_1 - C_6 -alkoxy, hydroxyl or C_1 - C_6 -alkoxycarbonyl;

 $R^7 \qquad \text{is C_4-C_6,-alkyl, C_3-C_6-alkenyl, C_3-C_6-alkynyl, C_3-C_6-alkynyl, C_3-C_6-alkynyl, C_3-C_6-alkenylcarbonyl, C_2-C_6-alkenylcarbonyl, C_2-C_6-alkenylcarbonyl, C_2-C_6-alkenylcarbonyl, C_3-C_6-alkenyloxycarbonyl, C_3-C_6-alkenyloxycarbonyl, C_3-C_6-alkynyloxycarbonyl, $(C_1$-C_2-alkylthio)carbonyl, $(C_1$-C_2-alkylaminocarbonyl, C_3-C_6-alkenylaminocarbonyl, C_3-C_6-alkenylaminocarbonyl, C_3-C_6-alkynylaminocarbonyl, N,N-di-(C_1-C_6-alkyl)aminocarbonyl, N,C_3-C_6-alkenyl)-N-(C_1-C_6-alkyl) aminocarbonyl , N-(C_3-C_6-alkynyl)-N-(C_1-C_6-alkyl) aminocarbonyl , N-(C_3-C_6-alkynyl)-N-(C_1-C_6-alkyl) aminocarbonyl ,$

N-(C_1 - C_6 -alkyl) aminocarbonyl , N-(C_3 - C_6 -alkenyl)-N-(C_1 - C_6 -alkoxy) aminocarbonyl , N-(C_3 - C_6 -alkynyl)-N-(C_1 - C_6 -alkoxy) aminocarbonyl, di-(C_1 - C_6 -alkyl)-aminothiocarbonyl, C_1 - C_6 -alkylcarbonyl- C_1 - C_6 -alkyl,

N-(C₁-C₆-alkoxy)-

 C_1 - C_6 -alkoxyimino- C_1 - C_6 -alkyl, N-(C₁-C₆-alkylamino) imino-C₁-C₆-alkyl or N,N-di-(C₁-C₆-alkylamino)imino-C₁-C₆-alkyl, it being possible for the above-mentioned alkyl, cycloalkyl and alkoxy radicals to be partially or fully halogenated and/or to carry one to three of the following groups: cyano, C_1 - C_2 -alkoxy, C_1 - C_2 -alkylthio, di- $(C_1$ - C_2 - alkyl)amino, C_1-C_4 -alkylcarbonyl, C_1-C_4 -alkoxycarbonyl, C_1-C_4 -alkoxy- C_1 - C_4 -alkoxycarbonyl, di- $(C_1$ - C_4 -alkyl)amino- C_1 - C_4 alkoxycarbonyl, hydroxycarbonyl, C₁-C₄-alkylaminocarbonyl, di-(C₁-C₄-alkyl)aminocarbonyl, aminocarbonyl, C₁-C₄alkylcarbonyloxy or C₃-C₆-cycloalkyl; phenyl, phenyl-C₁-C₂-alkyl, phenylcarbonyl-C₁-C₂-alkyl, phenylcarbonyl, phenoxycarbonyl, phenoxythiocarbonyl, phenoxy-C₁-C₆-alkylcarbonyl, phenylaminocarbonyl, N-(C₁-C₆-alkyl)-N-(phenyl)aminocarbonyl, or phenyl-C₂-C₆alkenylcarbonyl, it being possible for the phenyl radical of

the 10 last-mentioned substituents to be partially or fully halogenated and/or to carry one to three of the following radicals:

phenyl, phenyl- C_1 - C_6 -alkyl, heterocyclyl- C_1 - C_6 -alkyl, phenylcarbonyl- C_1 - C_6 -alkyl, heterocyclylcarbonyl- C_1 - C_6 -alkyl, phenylcarbonyl,

heterocyclylcarbonyl, phenoxycarbonyl, heterocyclyloxycarbonyl, phenoxy- C_1 - C_6 -alkylcarbonyl, heterocyclyloxy- C_1 - C_6 -alkylcarbonyl, phenylaminocarbonyl, N-(C_1 - C_6 -alkyl)-N-(phenyl)aminocarbonyl, heterocyclylaminocarbonyl, N-(C_1 - C_6 -alkyl)-N-(heterocyclyl)aminocarbonyl, phenyl- C_2 - C_6 -alkenylcarbonyl or heterocyclyl- C_2 - C_6 -alkenylcarbonyl, it being possible for the phenyl and the heterocyclyl radical of the 20 last-mentioned substituents to be partially or fully halogenated and/or to carry one to three of the following radicals:

nitro, cyano, C_1 - C_4 -alkyl, C_1 - C_4 -halogenalkyl, C_1 - C_4 -alkoxy or C_1 - C_4 -haloalkoxy;

 R^8, R^9 are C_1 - C_6 -alkyl, C_3 - C_6 -alkenyl, C_3 - C_6 -haloalkenyl, C_3 - C_6 -alkynyl, C_3 - C_6 -haloalkynyl, C_3 - C_6 -cycloalkyl, hydroxyl, C_1 - C_6 -alkoxy, amino, C_1 - C_6 -alkylamino, C_1 - C_6 -haloalkylamino, it being possible for the abovementioned alkyl, cycloalkyl and alkoxy radicals to be partially or fully halogenated and/or to carry one to three of the following groups: cyano, C_1 - C_4 -alkoxy, C_1 - C_4 -alkylthio, di- $(C_1$ - C_4 -alkyl)amino, C_1 - C_4 -alkylcarbonyl, C_1 - C_4 -alkoxycarbonyl, C_1 - C_4 -alkoxy- C_1 -

 C_4 -alkoxycarbonyl, di- $(C_1$ - C_4 -alkyl)amino- C_1 - C_4 -

alkoxycarbonyl,

hydroxycarbonyl, C_1 - C_4 -alkylaminocarbonyl, di- $(C_1$ - C_4 -alkyl)aminocarbonyl, aminocarbonyl, C_1 - C_4 -alkylcarbonyloxy or C_3 - C_6 -cycloalkyl;

phenyl, phenyl-C₁-C₆-alkyl, phenoxy, it being possible for the phenyl radical of the last-mentioned substituents to be partially or fully halogenated and/or to carry one to three of the following radicals:

phenyl, heterocyclyl, phenyl- C_1 - C_6 -alkyl, heterocyclyl- C_1 - C_6 -alkyl, phenoxy, heterocyclyloxy, it being possible for the phenyl and the heterocyclyl radical of the last-mentioned substituents to be partially or fully halogenated and/or to carry one to three of the following radicals: nitro, cyano, C_1 - C_4 -alkyl, C_1 - C_4 -haloalkyl, C_1 - C_4 -alkoxy or C_1 - C_4 -haloalkoxy;

 R^{10} is C_1 - C_6 -alkyl, C_3 - C_6 -alkenyl, C_3 - C_6 -haloalkenyl, C_3 - C_6 -alkynyl, C_3 - C_6 -haloalkynyl, C_3 - C_6 -cycloalkyl, hydroxyl, C_1 - C_6 -alkoxy, C_3 - C_6 -alkenyloxy, C_3 - C_6 -alkynyloxy, amino, C_1 - C_6 -alkylamino, di- $(C_1$ - C_6 -alkyl)amino or C_1 - C_6 -alkylcarbonylamino, where the abovementioned alkyl, cycloalkyl and alkoxy radicals may be partially or fully halogenated and/or may carry one to three radicals from the

following group:

cyano, C_1 - C_4 -alkoxy, C_1 - C_4 -alkylthio, di- $(C_1$ - C_4 -alkyl)amino, C_1 - C_4 -alkylcarbonyl, C_1 - C_4 -alkoxycarbonyl, C_1 - C_4 -alkoxycarbonyl, di- $(C_1$ - C_4 -alkyl)amino- C_1 - C_4 -alkoxycarbonyl, hydroxycarbonyl, C_1 - C_4 -alkylaminocarbonyl, di- $(C_1$ - C_4 -alkyl)aminocarbonyl, aminocarbonyl, C_1 - C_4 -alkyl)aminocarbonyl, aminocarbonyl, C_1 - C_4 -alkylcarbonyloxy or C_3 - C_6 -cycloalkyl; phenyl, or phenyl- C_1 - C_6 -alkyl, where the phenyl radical of the two last-mentioned substituents may be partially or fully halogenated and/or may carry one to three of the following

phenyl, heterocyclyl, phenyl-C₁-C₆-alkyl or heterocyclyl-C₁-C₆-alkyl, where the phenyl or heterocyclyl radical of the four last-mentioned substituents may be partially or fully halogenated and/or may carry one to three of the following radicals:

nitro, cyano, C_1 - C_4 -alkyl, C_1 - C_4 -haloalkyl, C_1 - C_4 -alkoxy or C_1 - C_4 -haloalkoxy;

 R^{11} , R^{12} are C_1 - C_6 -alkyl, C_3 - C_6 -alkenyl, C_3 - C_6 -alkynyl or C_1 - C_6 -alkylcarbonyl;

I is 0 to 6;

radicals:

- m is 2 to 4;
- n is 1 to 5;

- p is 2 to 5; and their agriculturally useful salts.
- 2. (currently amended) A cyclohexenonequinolinoyl derivative of the formula I as claimed in claim 1 where
 - is halogen, C₁-C₆-alkyl, C₁-C₆-haloalkyl, C₁-C₆-alkoxy, C₁-C₆-alkylthio, heterocyclyloxy or phenylthio, it being possible for the two last-mentioned radical radicals to be partially or fully halogenated and/or to carry one to two carry one to three of the substituents mentioned below:

 nitro, cyano, C₁-C₄-alkyl, C₁-C₄-haloalkyl, C₁-C₄-alkoxy or C₁-C₄-haloalkoxy;
 - is halogen, OR⁷, SR⁷, SOR⁸, SO₂R⁸, OSO₂R⁸, OPR⁸R⁹, OPOR⁸R⁹

 OPSR⁸R⁹, NR¹⁰R¹¹ or N-bonded heterocyclyl, which is unsubstituted or partially or fully halogenated and/or carries one to three of the following radicals:

nitro, cyano, C_1 - C_4 -alkyl, C_1 - C_4 -haloalkyl, C_1 - C_4 -alkoxy or C_1 - C_4 -haloalkoxy.

- 3. (currently amended) A cyclohexenonequinolinoyl derivative of the formula I as claimed in claim I, where
 - R⁵ is halogen, OR⁷, NR¹⁰R¹¹ or N-bonded heterocyclyl which is unsubstituted

ţ

or partially or fully halogenated and/or carries one to three of the following radicals:

nitro, cyano, C_1 - C_4 -alkyl, C_1 - C_4 -haloalkyl, C_1 - C_4 -alkoxy or C_1 - C_4 -haloalkoxy.

- 4. (previously presented) A cyclohexenonequinolinoyl derivative of the formula I as claimed in claim 1, where
 - R⁷ is C_1 - C_6 -alkyl, C_1 - C_{20} -alkylcarbonyl, C_1 - C_6 -alkoxycarbonyl, $(C_1$ - C_{20} -alkylthio)carbonyl, N_1 -di- $(C_1$ - C_6 -alkyl)aminocarbonyl, phenyl, phenylcarbonyl or phenoxy- C_1 - C_6 -alkylcarbonyl, it being possible for the phenyl radical of the three last-mentioned substituents to be partially or fully halogenated and/or to carry one to three of the following radicals: nitro, cyano, C_1 - C_4 -alkyl, C_1 - C_4 -haloalkyl, C_1 - C_4 -alkoxy or C_1 - C_4 -haloalkoxy;
 - R^{10} is C_1 - C_6 -alkyl or C_1 - C_6 -alkoxy;
 - R^{11} is C_1 - C_6 -alkyl.
- 5. (previously presented) A cyclohexenonequinolinoyl derivative of the formula I as claimed in claim 1, where
 - R⁶ is nitro, halogen, cyano, C₁-C₆-alkyl, C₁-C₆-haloalkyl, di-(C₁-

ľ

 $\label{eq:continuous_continuous$

or

two radicals , which are linked to the same carbon, together form an -O-(CH_2)_m-O-, -O-(CH_2)_m-S-, -S-(CH_2)_m-S-, -O-(CH_2)_n- or -S-(CH_2)_n chain which is unsubstituted or substituted by one to three radicals from the following group :

halogen, cyano, C₁-C₄-alkyl, C₁-C₄-haloalkyl or C₁-C₄-alkoxycarbonyl;

or

two radicals , which are linked to the same carbon, together form a $-(CH_2)_p$ chain which possibly is interrupted by oxygen or sulfur and which is unsubstituted or substituted by one to four radicals from the following group :

halogen, cyano, C_1 - C_4 -alkyl, C_1 - C_4 -haloalkyl or C_1 - C_4 -alkoxycarbonyl ; or

two radicals, which are linked to the same carbon, together with this carbon form a carbonyl group.

6. (previously presented) A process for preparing compounds of the formula I as claimed in claim 1 where R⁵ = halogen, which comprises reacting a cyclohexanedione derivative of the formula III,

$$(R^6)$$

where the variables R¹ to R³, and I are each as defined in claim 1, with a halogenating agent.

7. (previously presented) A process for preparing compounds of the formula I as claimed in claim 1 where R⁵ = OR⁷, OSO₂R⁸, OPR⁸R⁹, OPOR⁸R⁹ or OPSR⁸R⁹,

which comprises reacting a cyclohexanedione derivative of the formula III,

$$(R^6)$$

where the variables R^1 to R^3 , and I are each as defined in claim 1, with a compound of the formula $IV\alpha$, $IV\beta$, $IV\gamma$, $IV\delta$ or $IV\epsilon$,

$$L^1-R^7$$
 L^1-SO_2 R^8 $L^1-PR^8R^9$ $L^1-POR^8R^9$ $L^1-PSR^8R^9$ (IV α) (IV β) (IV γ) (IV δ) (IV ϵ)

where the variables R⁷ to R⁹ are each as defined in claim 1 and L¹ is a nucleophilically replaceable leaving group.

8. (currently amended) A process for preparing compounds of the formula I as claimed

in claim 1 where R⁵ = OR⁷, SR⁷, POR⁸R⁹, NR¹⁰R¹¹, ONR¹¹R¹², or N-linked heterocyclyl N-linked heterocyclyl or O-(N-linked heterocyclyl), which comprises reacting a compound of the formula I α (\equiv I where R⁵ = halogen, OSO₂R⁸),

$$(R^6) \xrightarrow{R^5} R^2$$
and/or
$$(R^6) \xrightarrow{R^5} R^2$$

$$(R^6) \xrightarrow{R^5} R^2$$

I where R⁵= halogen or OSO₂R8

where the variables R1 to R3, R6 and I are each as defined in claim 1, with a compound of the formula $V\alpha$, $V\beta$, $V\gamma$, $V\delta$, $V\epsilon$, $V\eta$, $V\vartheta$,

$$\begin{aligned} & HOR^7 & HSR^7 & HPOR^8R^9 & HNR^{10}R^{11} & HONR^{11}R^{12} \\ & (V\alpha) & (V\beta) & (V\gamma) & (V\delta) & (V\epsilon) \\ & & H(N\text{-linked} & & & \\ & & heterocyclyl) & & & & \\ & & V\eta & & & \\ \end{aligned}$$

where the variables R⁷ to R¹² are each as defined in claim 1, if appropriate

in the presence of a base.

9. (previously presented) A process for preparing compounds of the formula I as claimed in claim 1, where $R^5 = SOR^8$, SO_2R^8 , which comprises reacting a compound of the formula I β (\equiv I where $R^5 = SR^8$),

$$(\mathsf{R}^6) \overset{\mathsf{R}^3}{\longleftarrow} \mathsf{R}^2 \overset{\mathsf{R}^5}{\longleftarrow} \mathsf{R}^2$$
 and/or
$$(\mathsf{R}^6) \overset{\mathsf{R}^5}{\longleftarrow} \mathsf{R}^2$$

I where R5= SR8

where the variables R^1 to R^8 and I are each as defined in claim 1, with an oxidizing agent.

10. (previously presented) A composition, comprising a herbicidally effective amount of at least one cyclohexenonequinolinoyl derivative of the formula I or an agriculturally useful salt of formula I as claimed in claim 1 and auxiliaries which

are conventionally used for formulating crop protection agents.

- 11. (previously presented) A process for preparing a composition as claimed in claim 10, which comprises mixing a herbicidally effective amount of at least one cyclohexenonequinolinoyl derivative of the formula I or an agriculturally useful salt of formula I and auxiliaries which are conventionally used for formulating crop protection agents.
- 12. (previously presented) A method for controlling undesirable vegetation, which comprises allowing a herbicidally effective amount of at least one cyclohexenonequinolinoyl derivative of the formula I or an agriculturally useful salt of formula I as claimed in claim 1 to act on plants, their habitat and/or on seeds.
- 13. (canceled)
- 14. (currently amended) A cyclohexenonequinolinoyl derivative of the formula I

ŧ,

$$R^4$$
 R^2
 R^1

where:

R¹ is hydrogen, nitro, halogen, cyano, C_1 - C_6 -alkyl, C_1 - C_6 -haloalkyl, C_1 - C_6 -alkoxyiminomethyl, C_1 - C_6 -alkoxy, C_1 - C_6 -haloalkoxy, C_1 - C_6 -alkylthio, C_1 - C_6 -haloalkylthio, C_1 - C_6 -alkylsulfinyl, C_1 - C_6 -haloalkylsulfinyl, C_1 - C_6 -alkylsulfonyl, aminosulfonyl, C_1 - C_6 -alkylsulfonyl, aminosulfonyl, C_1 - C_6 -alkyl)aminosulfonyl,

N, N-di-(C_1 - C_6 -alkyl) aminosulfonyl, $N-(C_1-C_6-alkylsulfonyl)amino, \\ N-(C_1-C_6-haloalkylsulfonyl)amino, \\ N-(C_1-C_6-alkyl)-N-(C_1-C_6-alkylsulfonyl)amino,$

N-(C_1 - C_6 --alkyl)-N-(C_1 - C_6 -haloalkylsulfonyl)amino, phenoxy, heterocyclyloxy, or phenylthio or heterocyclylthio, it being possible for the two four last-mentioned radicals to be partially or fully halogenated and/or to carry one to two one to three of the following substituents:

nitro, cyano, C₁-C₄-alkyl, C₁-C₄-haloalkyl,

 C_1 - C_4 -alkoxy or C_1 - C_4 -haloalkoxy;

 R^2 , R^3 are hydrogen, C_1 - C_6 -alkyl, C_1 - C_6 -haloalkyl or halogen;

R⁴ is a compound IIa

$$(\mathsf{R}^6) \xrightarrow[\mathsf{R}^5]{} (\mathsf{R}^6) \xrightarrow[\mathsf{Ilb}]{} \mathsf{Ilb}$$

where

R⁵ is halogen, OR⁷, SR⁷, SOR⁸, SO₂R⁸, OSO₂R⁸, POR⁸R⁹,

OPR⁸R⁹, OPOR⁸R⁹, OPSR⁸R⁹, NR¹⁰R¹¹, ONR¹¹R¹², N-linked heterocyclyl or O-(N-linked heterocyclyl), it being possible for the heterocyclyl radical of the two last-mentioned substituents to be partially or fully halogenated and/or to carry one to three of the following radicals:

nitro, cyano, C₁-C₄-alkyl, C₁-C₄-haloalkyl, C₁-C₄-alkoxy or C₁-C₄-haloalkoxy;

R⁶ is nitro, halogen, cyano, C₁-C₆-alkyl,

C₁-C₆-haloalkyl, di-(C₁-C₆-alkoxy)methyl,

di-(C₁-C₆-alkylthio)methyl,

(C₁-C₆-alkoxy)(C₁-C₆-alkylthio)methyl, hydroxyl,

 C_1 - C_6 -alkoxy, C_1 - C_6 -haloalkoxy,

C₁-C₆-alkoxycarbonyloxy, C₁-C₆-alkylthio,

 C_1 - C_6 -haloalkylthio, C_1 - C_6 -alkylsulfinyl,

 C_1 - C_6 -haloalkylsulfinyl, C_1 - C_6 -alkylsulfonyl,

 C_1 - C_6 -haloalkylsulfonyl, C_1 - C_6 -alkylcarbonyl,

 C_1 - C_6 -haloalkylcarbonyl, C_1 - C_6 -alkoxycarbonyl or

C₁-C₆-haloalkoxycarbonyl;

or

two radicals , which are linked to the same carbon, together form an -O-(CH_2)_m-O-, -O-(CH_2)_m-S-, -S-(CH_2)_m-S-, -

 $O-(CH_2)_n$ - or $-S-(CH_2)_n$ chain which is unsubstituted or substituted by one to three radicals from the following group: halogen, cyano, C_1-C_4 -alkyl, C_1-C_4 -haloalkyl or C_1-C_4 -alkoxycarbonyl;

or

ť

two radicals , which are linked to the same carbon, together form a -(CH_2)_p chain which possibly is interrupted by oxygen or sulfur and/or is unsubstituted or substituted by one to four radicals from the following group: halogen, cyano, C_1 - C_4 -alkyl, C_1 - C_4 -haloalkyl or C_1 - C_4 -alkoxycarbonyl;

or

two radicals , which are linked to the same carbon, together form a methylidene group which is unsubstituted or substituted by one or two radicals from the following group: halogen, hydroxyl, formyl, cyano, C_1 - C_6 -alkyl, C_1 - C_6 -haloalkyl, C_1 - C_6 -alkoxy, C_1 - C_6 -haloalkoxy, C_1 - C_6 -alkylthio, C_1 - C_6 -haloalkylthio, C_1 - C_6 -haloalkylthio, C_1 - C_6 -alkylsulfinyl, C_1 - C_6 -alkylsulfonyl or C_1 - C_6 -haloalkylsulfonyl;

or

two radicals, which are linked to the same carbon,

together with this carbon form a carbonyl group;

or

two radicals , which are linked to different carbons, together form a $-(CH_2)_n$ chain which is unsubstituted or substituted by one to three radicals from the following group: halogen, C_1 - C_6 -alkyl, C_1 - C_6 -alkoxy, hydroxyl or C_1 - C_6 -alkoxycarbonyl;

 $R^7 \qquad \text{is C_1-C_6,-alkyl, C_3-C_6-alkenyl, C_3-C_6-haloalkenyl,} \\ C_3-C_6-alkynyl, C_3-C_6-haloalkynyl, C_3-C_6-cyloalkyl,} \\ C_1-C_{20}-alkylcarbonyl, C_2-C_6-alkenylcarbonyl,} \\ C_2-C_6-alkynylcarbonyl, C_3-C_6-cyloalkylcarbonyl,} \\ C_1-C_6-alkoxycarbonyl, C_3-C_6-alkenyloxycarbonyl,} \\ C_3-C_6-alkynyloxycarbonyl,} \\ (C_1-C_{20}-alkylthio)carbonyl,} \\ C_3-C_6-alkylaminocarbonyl,} \\ C_3-C_6-alkenylaminocarbonyl,} \\ C_3-C_6-alkynylaminocarbonyl,} \\ N,N-di-(C_1-C_6-alkyl)aminocarbonyl,} \\ N-(C_3-C_6-alkenyl)-N-(C_1-C_6-alkyl) aminocarbonyl,} \\ N-(C_3-C_6-alkenyl)-N-(C_1-C_6-alkyl) aminocarbonyl,} \\ N-(C_3-C_6-alkynyl)-N-(C_1-C_6-alkyl) aminocarbonyl,} \\ N-(C_3-C_6-alkynyl)-N-(C_3-C_6-alkyl)-N-(C_3-C_6-alkyl)-N-(C_3-C_6-alkyl)-N-(C_3-C_6-alkyl)-N-(C_3-C_6-alkyl)-N-(C_3-C_6-alkyl)-N-(C_3-C_6-alkyl)-N-(C_3-C_6-alkyl)-N-(C_3-C_6-alkyl)-N-(C_3-$

4. 6.6

 $N-(C_1-C_6-alkoxy)-$

N-(C_1 - C_6 -alkyl) aminocarbonyl, N-(C_3 - C_6 -alkenyl)-

 $N-(C_1-C_6-alkoxy)$ aminocarbonyl, $N-(C_3-C_6-alkynyl)$ -

 $N-(C_1-C_6-alkoxy)$ aminocarbonyl, di- $(C_1-C_6-alkyl)$ -

aminothiocarbonyl, C₁-C₆-alkylcarbonyl-C₁-C₆-alkyl,

C₁-C₆-alkoxyimino-C₁-C₆-alkyl,

N-(C_1 - C_6 -alkylamino) imino- C_1 - C_6 -alkyl or

N,N-di-(C_1 - C_6 -alkylamino)imino- C_1 - C_6 -alkyl, it being possible for the above-mentioned alkyl, cycloalkyl and alkoxy radicals to be partially or fully halogenated and/or to carry one to three of the following groups: cyano, C_1 - C_4 -alkoxy, C_1 - C_4 -alkylthio, di-(C_1 - C_4 -alkyl) amino, C_1 - C_4 -alkylcarbonyl, C_1 - C_4 -alkoxycarbonyl, C_1 - C_4 -alkoxycarbonyl, di-(C_1 - C_4 -alkoxycarbonyl, hydroxycarbonyl, C_1 - C_4 -alkylaminocarbonyl, di-(C_1 - C_4 -alkyl)aminocarbonyl, aminocarbonyl, C_1 - C_4 -alkylcarbonyloxy or C_3 - C_6 -cycloalkyl;

phenyl, phenyl- C_1 - C_6 -alkyl, phenylcarbonyl- C_1 - C_6 -alkyl, phenylcarbonyl, phenoxycarbonyl, phenoxythiocarbonyl, phenoxy- C_1 - C_6 -alkylcarbonyl, phenylaminocarbonyl, N-(C_1 - C_6 -alkyl)-N-(phenyl)aminocarbonyl, or phenyl- C_2 - C_6 -alkenylcarbonyl, it being possible for the phenyl radical of the 10 last-mentioned substituents to be partially or fully halogenated and/or to carry one to three of the following radicals:

£ , 4 15

phenyl, heterocyclyl, phenyl- C_1 - C_6 -alkyl, heterocyclyl- C_1 - C_6 -alkyl, phenylcarbonyl- C_1 - C_6 -alkyl, heterocyclylcarbonyl- C_1 - C_6 -alkyl, phenylcarbonyl, heterocyclylcarbonyl, phenoxycarbonyl, heterocyclyloxycarbonyl, phenoxythiocarbonyl, heterocyclyloxythiocarbonyl, phenoxy- C_1 - C_6 -alkylcarbonyl, heterocyclyloxy- C_1 - C_6 -alkylcarbonyl, phenylaminocarbonyl, N- $(C_1$ - C_6 -alkyl)-N-(phenyl)aminocarbonyl, heterocyclylaminocarbonyl, N- $(C_1$ - C_6 -alkyl)-N-(heterocyclyl)aminocarbonyl, phenyl- C_2 - C_6 -alkenylcarbonyl or heterocyclyl- C_2 - C_6 -alkenylcarbonyl, it being possible for the phenyl and the heterocyclyl radical of the 20 last-mentioned substituents to be partially or fully halogenated and/or to carry one to three of the following radicals:

nitro, cyano, C_1 - C_4 -alkyl, C_1 - C_4 -halogenalkyl, C_1 - C_4 -alkoxy or C_1 - C_4 -haloalkoxy;

 R^8 , R^9 are C_1 - C_6 -alkyl, C_3 - C_6 -alkenyl, C_3 - C_6 -haloalkenyl, C_3 - C_6 -alkynyl, C_3 - C_6 -cycloalkyl, hydroxyl, C_1 - C_6 -alkoxy, amino, C_1 - C_6 -alkylamino, C_1 - C_6 -haloalkylamino, di- $(C_1$ - C_6 -alkyl) amino or di- $(C_1$ - C_6 -haloalkyl) amino, it being possible for the abovementioned alkyl, cycloalkyl and alkoxy radicals to be partially or fully halogenated and/or to carry one to three of the following groups:

C₄-haloalkoxy;

J. 7 7

alkylcarbonyl, C_1 - C_4 -alkoxycarbonyl, C_1 - C_4 -alkoxy- C_1 - C_4 -alkoxycarbonyl, di- $(C_1$ - C_4 -alkyl)amino- C_1 - C_4 -alkoxycarbonyl, di- $(C_1$ - C_4 -alkyl)aminocarbonyl, aminocarbonyl, di- $(C_1$ - C_4 -alkylcarbonyloxy or C_3 - C_6 -cycloalkyl; phenyl, phenyl- C_1 - C_6 -alkyl, phenoxy, it being possible for the phenyl radical of the last-mentioned substituents to be partially or fully halogenated and/or to carry one to three of the following radicals: phenyl, heterocyclyl, phenyl- C_1 - C_6 -alkyl, heterocyclyl- C_1 - C_6 -alkyl, phenoxy, heterocyclyloxy, it being possible for the phenyl and the heterocyclyl radical of the last-mentioned substituents to be partially or fully halogenated and/or to carry one to three of the following radicals: nitro, cyano, C_1 - C_4 -alkyl, C_1 - C_4 -haloalkyl, C_1 - C_4 -alkoxy or C_1 -

cyano, C_1 - C_4 -alkoxy, C_1 - C_4 -alkylthio, di- $(C_1$ - C_4 - alkyl)amino, C_1 - C_4 -

R¹⁰ is C_1 - C_6 -alkyl, C_3 - C_6 -alkenyl, C_3 - C_6 -haloalkenyl, C_3 - C_6 -alkynyl, C_3 - C_6 -haloalkynyl, C_3 - C_6 -cycloalkyl, hydroxyl, C_1 - C_6 -alkoxy, C_3 - C_6 -alkenyloxy, C_3 - C_6 -alkynyloxy, amino, C_1 - C_6 -alkylamino, di-(C_1 - C_6 -alkyl)amino or C_1 - C_6 -alkylcarbonylamino, it being possible for the abovementioned alkyl, cycloalkyl and alkoxy radicals to be

18. 17. A.

partially or fully halogenated and/or to carry one to three radicals from the following group:

cyano, C₁-C₄-alkoxy, C₁-C₄-alkylthio, di-(C₁-C₄-alkyl)amino,

C₁-C₄-alkylcarbonyl, C₁-C₄-alkoxycarbonyl, C₁-C₄-alkoxy-C₁-

C₄-alkoxycarbonyl, di-(C₁-C₄-alkyl)amino-C₁-C₄-

alkoxycarbonyl, hydroxycarbonyl, C₁-C₄-alkylaminocarbonyl,

di-(C₁-C₄-alkyl)aminocarbonyl, aminocarbonyl, C₁-C₄-

alkylcarbonyloxy or C₃-C₆-cycloalkyl;

phenyl, or phenyl- C_1 - C_6 -alkyl, it being possible for the phenyl radical of the two last-mentioned substituents to be partially or fully halogenated and/or to carry one to three of the following radicals:

phenyl, heterocyclyl, phenyl-C₁-C₆-alkyl or heterocyclyl-C₁-C₆-alkyl, it being possible for the phenyl or heterocyclyl radical of the four last-mentioned substituents to be partially or fully halogenated and/or to carry one to three of the following radicals:

nitro, cyano, C_1 - C_4 -alkyl, C_1 - C_4 -haloalkyl, C_1 - C_4 -alkoxy or C_1 - C_4 -haloalkoxy;

 R^{11} , R^{12} are C_1 - C_6 -alkyl, C_3 - C_6 -alkenyl, C_3 - C_6 -alkynyl or C_1 - C_6 -alkylcarbonyl;

l is 0 to 6:

m is 2 to 4:

- n is 1 to 5;
- p is 2 to 5;

and their agriculturally useful salts.

- 15. (currently amended) A cyclohexenonequinolinoyl derivative of the formula I as claimed in claim 14, where
 - is halogen, C₁-C₆-alkyl, C₁-C₆-haloalkyl, C₁-C₆-alkoxy, C₁-C₆-alkylthio, heterocyclyloxy or phenylthio, it being possible for the two last-mentioned radical radicals to be partially or fully halogenated and/or to carry one to two one to three of the substituents mentioned below:

 nitro, cyano, C₁-C₄-alkyl, C₁-C₄-haloalkyl, C₁-C₄-alkoxy or C₁-C₄-haloalkoxy;
 - is halogen, OR⁷, SR⁷, SOR⁸, SO₂R⁸, OSO₂R⁸, OPR⁸R⁹, OPOR⁸R⁹

 OPSR⁸R⁹, NR¹⁰R¹¹ or N-bonded heterocyclyl which is unsubstituted or partially or fully halogenated and/or carries one to three of the following radicals:

nitro, cyano, C_{\uparrow} - C_{\sharp} -alkyl, C_{\uparrow} - C_{\sharp} -haloalkyl, C_{\uparrow} - C_{\sharp} -alkoxy or C_{\uparrow} - C_{\sharp} -haloalkoxy .

16. (currently amended) A cyclohexenonequinolinoyl derivative of the formula I as claimed in claim 14, where

haloalkoxy.

17

- is halogen, OR⁷, NR¹⁰R¹¹ or N-bonded heterocyclyl which is unsubstituted or partially or fully halogenated and/or carries one to three of the following radicals:

 nitro, cyano, C₁-C₄-alkyl, C₁-C₄-haloalkyl, C₁-C₄-alkoxy or C₁-C₄-
- 17. (currently amended) A cyclohexenonequinolinoyl derivative of the formula I as claimed in claim 14, where

 - R^{10} is C_1 - C_6 -alkyl or C_1 - C_6 -alkoxy;
 - R^{11} is C_1 - C_6 -alkyl.
- 18. (previously presented) A process for preparing compounds of the formula I as claimed in claim 14 where R⁵ = halogen, which comprises reacting a

cyclohexanedione derivative of the formula III,

$$(R^6)$$

where the variables R^1 to R^3 , and I are each as defined in claim 14, with a halogenating agent.

19. (previously presented) A process for preparing compounds of the formula I as claimed in claim 14 where R⁵ = OR⁷, OSO₂R⁸, OPR⁸R⁹, OPOR⁸R⁹ or OPSR⁸R⁹, which comprises reacting a cyclohexanedione derivative of the formula III,

$$(R^6)$$

where the variables R^1 to R^3 , and I are each as defined in claim 14, with a compound of the formula $IV\alpha$, $IV\beta$, $IV\gamma$, $Iv\delta$ or $IV\epsilon$,

$$L^{1}-R^{7}$$
 $L^{1}-SO_{2}$ R^{8} $L^{1}-PR^{8}R^{9}$ $L^{1}-POR^{8}R^{9}$ $L^{1}-PSR^{8}R^{9}$ (IV α) (IV β) (IV γ) (IV δ) (IV ϵ)

where the variables R^7 to R^9 are each as defined in claim 14 and L^1 is a nucleophilically replaceable leaving group.

20. (currently amended) A process for preparing compounds of the formula I as claimed in claim 14 where R⁵ = OR⁷, SR⁷, POR⁸R⁹, NR¹⁰R¹¹, ONR¹¹R¹², or N-linked

heterocyclyl N-linked heterocyclyl or O-(N-linked heterocyclyl), which comprises reacting a compound of the formula I α (\equiv I where R⁵ = halogen, OSO₂R⁸),

$$(\mathsf{R}^6) \xrightarrow{\mathsf{R}^3} \mathsf{R}^2$$
 and/or
$$(\mathsf{R}^6) \xrightarrow{\mathsf{R}^5} \mathsf{R}^2$$

I where R⁵= halogen or OSO₂R⁸

HOR ⁷	HSR ⁷	HPOR [®] R [®] H	NR ¹⁰ R ¹¹	HONR ¹¹ R ¹²
(Va)	(Vβ)	(Vγ)	(Vδ)	(Vε)
H(N-linked	H(ON-linked			
heterocyclyl	l) heterocyclyl)			
Vη	₩			

where the variables R7 to R12 are each as defined in claim 14, if

appropriate in the presence of a base.

21. (previously presented) A process for preparing compounds of the formula I as claimed in claim 14 where $R^5 = SOR^8$, SO_2R^8 , which comprises reacting a compound of the formula I β (\equiv I where $R^5 = SR^8$),

$$(\mathsf{R}^6) \overset{\mathsf{R}^3}{\longleftarrow} \mathsf{R}^2$$
 and/or
$$(\mathsf{R}^6) \overset{\mathsf{R}^5}{\longleftarrow} \mathsf{R}^1$$

I where R5= SR8

where the variables R¹ to R⁵, R⁵, R³ and I are each as defined in claim 14, with an oxidizing agent.

22. (currently amended) A composition, comprising a herbicidally effective amount of at

least one cyclohexenonequinolinoyl derivative of the formula I or an agriculturally useful salt of formula I as claimed in claim 14 and conventional crop protection formulation auxiliaries which are conventionally used for formulating crop protection agents.

- 23. (currently amended) A process for preparing a composition as claimed in claim 22, which comprises mixing a herbicidally effective amount of at least one cyclohexenonequinolinoyl derivative of the formula I or an agriculturally useful salt of formula I and conventional crop protection formulation auxiliaries which are conventionally used for formulating crop protection agents.
- 24. (previously presented) A method for controlling undesirable vegetation, which comprises allowing a herbicidally effective amount of at least one cyclohexenonequinolinoyl derivative of the formula I or an agriculturally useful salt of formula I as claimed in claim 14 to act on plants, their habitat and/or on seeds.

25. (canceled)